

Confinement and monopole condensation: some properties of the disorder parameter.

A. Di Giacomo^{a*} and G. Paffuti^a

^aDip. Fisica Università and INFN Pisa
Via Buonarroti 2, 56125 Pisa

1. Introduction

We consider a $U(1)$ gauge theory with a conserved magnetic current $j_\mu^M = \partial^\nu F_{\mu\nu}^*$. The corresponding magnetic $U(1)$ symmetry can either be Wigner or broken à la Higgs. In the first case (Wigner) the Hilbert space consists of superselected sectors with definite magnetic charge. In the second case (Higgs) at least one magnetically charged operator μ exists, with $\langle\mu\rangle \neq 0$. $\langle\mu\rangle$ is the order parameter. The free energy density (effective Lagrangean), is uniquely determined by symmetry and dimensional arguments[2] to be

$$\mathcal{L} = (D_\rho \langle\mu\rangle)^* D_\rho \langle\mu\rangle - \frac{1}{4} F^{\mu\nu} F_{\mu\nu} - V(\langle\mu\rangle) \quad (1)$$

Here $D_\rho = \partial_\rho - i q_M \tilde{A}_\rho$ is the dual covariant derivative, $\tilde{A}_\rho$ the dual vector potential, $V(\langle\mu\rangle)$ the usual quartic potential. If V has a non trivial minimum, $\langle\mu\rangle \neq 0$, the system is a dual superconductor. In the usual non compact formulation $j_\mu^M \equiv 0$ (Bianchi identities). On a lattice the theory is compact and a non zero magnetic

current can be defined[3]. In QCD $U(1)$ is defined by abelian projection[4]. A magnetically charged operator μ has been constructed[1] and $\langle\mu\rangle$ has been used as a disorder parameter for detecting dual superconductivity. The results of this investigation are the following[5]: 1) The confined phase of quenched QCD has $\langle\mu\rangle \neq 0$, and behaves as a dual superconductor in all the abelian projections[1]. 2) In the deconfined phase $\langle\mu\rangle = 0$ and superselected magnetic sectors exist. 3) In the vicinity of the transition $\langle\mu\rangle \propto_{T \rightarrow T_c^-} \left(1 - \frac{T}{T_c}\right)^\delta$. δ is independent of the abelian projection, and is equal to the analogous index of the dual Polyakov line[6]. 4) A similar behaviour is found in the presence of dynamical quarks. These results are obtained by properly performing the infinite volume limit, by use of finite size scaling techniques[1,5,6], and provide basic information on the dual structure of QCD. The questions we want to address here are the following
1) Does $\langle\mu\rangle \neq 0$ contradict the so called Elitzur's theorem[8]? 2) How precisely is the abelian projection definable on the lattice?

*Speaker at the Conference. Partially supported by MURST and by EC, FMRX-CT97-0122

2. The order parameter of a superconductor.

The ground state of a superconductor is a superposition of states with different charges[9], and the order parameter $\langle\varphi\rangle$ is the v.e.v. of a charged operator φ . **Theorem** (continuum version of ref.[8]) If $\varphi(x)$ is a local charged operator, in a gauge invariant formulation (no gauge fixing), $|0\rangle$ is gauge invariant and hence

$$\langle 0|\varphi(x)|0\rangle = e^{i\Lambda(x)}\langle 0|\varphi(x)|0\rangle \quad \forall \Lambda(x)$$

or $\langle 0|\varphi(x)|0\rangle = 0$. Does the existence of a superconductor violate gauge invariance? The answer is of course no. In the usual perturbative treatment of the Higgs phenomenon a gauge is fixed, e.g. the unitary gauge[10], and the theorem is eluded. In a gauge invariant formulation, like lattice, the way out is to define gauge invariant charged operators[11], $\tilde{\varphi}(x)$, as follows

$$\tilde{\varphi}(x) = \varphi(x)e^{i(A,h)} \quad (2)$$

where

$$(A, h) = \int d^4y A_\mu(y) h_\mu(y - x)$$

$\partial^\mu h_\mu(z) = \delta^4(z)$. Under a generic gauge transformation U_Λ , with $\Lambda(x) \rightarrow 0$ as $|x| \rightarrow \infty$, $(A, h) \rightarrow (A, h) - \Lambda(x)$, $\varphi(x) \rightarrow e^{i\Lambda(x)}\varphi(x)$ and hence $\tilde{\varphi}(x) \rightarrow \tilde{\varphi}(x)$. $\tilde{\varphi}(x)$ is gauge invariant. However, under a global transformation (with $\partial_\mu \Lambda = 0$) $A_\mu \rightarrow A_\mu$ and $\tilde{\varphi}(x) \rightarrow e^{i\Lambda}\tilde{\varphi}(x)$ like any charged operator. $\tilde{\varphi}(x)$ is charged and gauge invariant. It is non local, but by a judicious choice of h_μ , it obeys cluster property

$$G(x) \equiv \langle \tilde{\varphi}(x)^\dagger \tilde{\varphi}(0) \rangle \underset{|x| \rightarrow \infty}{\simeq} A e^{-M|x|} + |\langle \varphi \rangle|^2 \quad (3)$$

and defines the order parameter $\langle\varphi\rangle$ by the asymptotic behaviour. Possible choices for h_μ depend on the dimension d of its support. For $d = 1$ $h^\mu = \delta_0^\mu \theta(x_0 - y_0) \delta^3(\vec{x} - \vec{y})$ and $\exp(i(A, h)) = \exp(i \int_{-\infty}^{x_0} A_0(y_0, \vec{x}) dy_0)$ is a parallel transport from $-\infty$ along time axis (Mandelstam string). The string can be put on any path C going to infinity and $\tilde{\varphi}(x) = \varphi(x) \exp(i \int_C A_\mu dx^\mu)$. For $d = 3$ $h^\mu = (0, \frac{1}{4\pi} \frac{\vec{x} - \vec{y}}{|\vec{x} - \vec{y}|^3})$ (Dirac choice). For a lattice version of the Higgs model

$$\mathcal{L} = \frac{\beta}{2} - a^2 \cos(d\theta - eA) \quad (4)$$

it can be proved that

- 1) At sufficiently small a (Coulomb phase) $|G(x)| \leq_{|x| \rightarrow \infty} C \exp(-\rho|x|)$
- 2) At sufficiently large a and small e (Higgs phase) $|G(x)| \geq a^2$ as $|x| \rightarrow \infty$, provided (h, h) is uniformly bounded in the limit $V \rightarrow \infty$. This is true for the Dirac choice of h^μ ($d = 3$) but not for the string ($d = 1$).
- 3) The Hilbert space enlarged by the correlators of $\tilde{\varphi}$ obeys Osterwalder-Schrader positivity, and admits a decomposition in superselected sectors in the Coulomb phase.

3. Dual superconductivity.

For compact $U(1)$ the operator μ which creates a monopole is defined as[12]

$$\mu(x_0, \vec{x}) = e^{\beta \sum_{\vec{n}} (\cos[\theta_{0i}(x_0, \vec{n}) - b_i(\vec{x} - \vec{n})] - \cos \theta_{0i}(x_0, \vec{n}))}$$

where $b_i(\vec{x} - \vec{y})$ is the vector potential at $\vec{y}$ of a monopole sitting at $\vec{x}$. Theorems

- 1) μ carries magnetic charge[12].
- 2) μ is gauge invariant ($\theta_{0i} = F_{0i}$).
- 3) μ is Dirac

like[13,14]. 4)The correlations functions $\langle \mu(x) \bar{\mu}(0) \rangle$ are equal to $\langle \tilde{\varphi}(x) \tilde{\varphi}^\dagger(0) \rangle$ of the Higgs model with the change $\beta_H \rightarrow \beta = \frac{1}{4\pi^2 \beta_H}$, as required by duality. In conclusion μ is a gauge invariant charged operator and $\langle \mu \rangle$ a legitimate disorder parameter for dual superconductivity.

4. About the abelian projection.

The abelian projection is a gauge transformation which diagonalizes an operator $\Phi(x) = \sum \Phi^a T^a$ in the adjoint representation. In this gauge the generic link on the lattice $U_\mu(n)$ can be written in the form

$$U_\mu(n) = V_\nu(n) C_\mu(n) \quad (5)$$

where $C_\mu(n)$ is an exponent of the diagonal generators (photons), $V_\mu(n)$ of the off diagonal (charged) generators. C_μ is uniquely defined. Also the form

$$U_\mu(n) = C_\mu(n) V'_\mu(n)$$

is possible, with $V'_\mu = C_\mu^\dagger(n) V_\mu C_\mu(n)$, again a charged operator. For the plaquette $\Pi_{\mu\nu} = U_\mu(n) U_\nu(n + \hat{\mu}) U_\mu^\dagger(n + \hat{\nu}) U_\nu^\dagger(n)$ the abelian projection is not uniquely defined. It can be defined, e.g., as the projected part of $\Pi_{\mu\nu}$ by a procedure similar to that leading to eq.(1), or, as usually done, as the plaquette constructed with the abelian links $\Pi_{\mu\nu}^0 = C_\mu(n) C_\nu(n + \hat{\mu}) C_\mu^\dagger(n + \hat{\nu}) C_\nu^\dagger(n)$. By use of eq.(7)

$$\Pi_{\mu\nu} = V_\mu(n) V'_\nu(n + \hat{\mu}) (V''_\mu)^\dagger(n + \hat{\nu}) (V'''_\nu)^\dagger(n) \Pi_{\mu\nu}^0 \quad (6)$$

The V 's in eq.(8) are separately charged, but their product contains at the exponent terms coming from the commutators

in the Baker-hausdorff formula, which are $\mathcal{O}(a^2)$ and belong to the diagonal subspace of the algebra. The abelian projection on the lattice is undefined by terms $\mathcal{O}(a^2)$.

REFERENCES

1. A. Di Giacomo, B. Lucini, L. Montesi and G. Paffuti *Phys. Rev.* **61** 034504,034505 (2000).
2. V.L. Ginzburg, L.D. Landau, *Zh. Eksp. Teor. Fiz.* **24**, 1064, (1950).
3. T.A. Degrand, D. Toussaint, *Phys. Rev.* **D22**, 2478, (1980).
4. G. 't Hooft, *Nucl. Phys.* **B 190**, 455 (1981).
5. J.M. Carmona, M. D'Elia, A. Di Giacomo, B. Lucini, hep-lat 0103005, to appear in *Phys. Rev. D*.
6. L. Del Debbio, A. Di Giacomo, B. Lucini, *Nucl. Phys.* **B 594**, 287 (2001); *Phys. Lett.* **B500**, 326 (2001).
7. J.M.Carmona, M.D'Elia, A.Di Giacomo, B.Lucini, these proceedings.
8. S. Elitzur, *Phys. Rev.* **D 12**, 3978 (1975).
9. N.N. Bogoliubov, *Nuovo Cimento* **7**, (1958) 794; J.G. Valatin, *Nuovo Cimento* **7**, (1958) 843.
10. S. Weinberg, *The Quantum Theory of Fields*, II, Cambridge University Press, 1996.
11. P.A.M. Dirac, *Can. Journ. of Phys.* **33**, 650 (1955).
12. A. Di Giacomo, G. Paffuti, *Phys. Rev.* **D56**, 6816 (1997).
13. J. Frölich, P.A. Marchetti, *Phys. Rev.* **D 64**, 014505 (2001).
14. V.Cirigliano, G.Paffuti, *Comm. Math. Phys.* **200**, 381, (1999)